

MCQS^{Q&As}

Multiple-choice questions for general practitioner (GP) Doctor

Pass Test Prep MCQS Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.leads4pass.com/mcqs.html>

100% Passing Guarantee
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers



QUESTION 1

An elderly man comes to the emergency department with fever, headache, a stiff neck, and photophobia. He is HIV positive with

What is the best initial therapy?

- A. Cefoxitin and mefloquine
- B. Fluconazole
- C. Ceftriaxone, ampicillin, and vancomycin

Correct Answer: C

Listeria monocytogenes is a cause of meningitis that is not adequately treated by any form of cephalosporin. Ampicillin is added to the usual regimen of ceftriaxone and vancomycin to cover Listeria. This cannot be fungal meningitis, because the CSF is characterized exclusively by a high number of neutrophils; neutrophils are not consistent with fungal meningitis.

QUESTION 2

A 32-year-old female who has had 3 previous miscarriages in the 1st trimester now comes with vaginal bleeding at 8 weeks. US reveals a viable fetus.

What would be the most appropriate definitive management?

- A. No treatment
- B. Bed rest 2 weeks
- C. Admit
- D. Aspirin

Correct Answer: D

QUESTION 3

A male presented with headache, tinnitus and nausea thinking that he has a brain tumor. He had just secured a job in a prestigious company and he thinks that he might not meet its standards. CNS exam, CT, all within normal.

What is the diagnosis?

- A. Conversion reaction
- B. Generalized Anxiety disorder
- C. Panic attack
- D. Hypochondriasis

Correct Answer: D

Diagnostic criteria for hypochondriasis include the following: The patient has a preoccupying fear of having a serious disease. The preoccupation persists despite appropriate medical evaluation and reassurance. The belief is not of delusional intensity (as in delusional disorder, somatic type) and is not Restricted to a concern about appearance (as in persons with BDD). The preoccupation causes clinically significant distress or impairment.

QUESTION 4

A person after oral sex diagnose with neisseria gonorrhoeae infection. Which is true regarding scenario?

- A. Treat his friends
- B. Treat his sexual partner
- C. Treat his family members

Correct Answer: B

QUESTION 5

A 73-year-old man who is recovering from surgery on the left carotid artery in his neck. He has slurred speech of his tongue, the tip deviated to the left.

What is the single most appropriate option?

- A. Accessory nerve
- B. Hypoglossal nerve
- C. Glossopharyngeal nerve
- D. Facial nerve
- E. Vagus nerve

Correct Answer: B

Testing function of the nerve is performed by asked the subject to stick their ton out. If there is a loss of innervation to one side, the tongue will curve toward the affected side, due to unopposed the opposite genioglossus muscle. If this is the result of. A lower motor neuron lesion, the tongue will be curve damaged side, combined with the presence of fasciculations or atrophy. However, if the deficit is caused by a neuron lesion, the tongue will be curved away from the side of the cortical damage, without the presence of fa or atrophy.

[Latest MCQS Dumps](#)

[MCQS Exam Questions](#)

[MCQS Braindumps](#)