

## MCQS<sup>Q&As</sup>

Multiple-choice questions for general practitioner (GP) Doctor

### Pass Test Prep MCQS Exam with 100% Guarantee

Free Download Real Questions & Answers **PDF** and **VCE** file from:

<https://www.leads4pass.com/mcqs.html>

100% Passing Guarantee  
100% Money Back Assurance

Following Questions and Answers are all new published by Test Prep  
Official Exam Center

- ⚙️ **Instant Download** After Purchase
- ⚙️ **100% Money Back** Guarantee
- ⚙️ **365 Days** Free Update
- ⚙️ **800,000+** Satisfied Customers



**QUESTION 1**

A 43-years-old man is brought to the emergency department after a motor vehicle accident involving a car. He mentioned that he is having headache and dizziness. During his overnight admission for observation, his serum sodium level rises to 151 meq/L.

All of the following tests are indicated EXCEPT:

- A. Measurement of morning cortisol level
- B. Measurement of response to desmopressin
- C. MRI scan of the head
- D. Overnight dehydration test

Correct Answer: A

---

**QUESTION 2**

Which of the following is not a degenerative corneal disease?

- A. Keratokonus
- B. Marginal dystrophy
- C. Fatty/lipid degeneration
- D. Mooren's ulcer
- E. Band keratopathy

Correct Answer: D

All other options are degenerative disease of cornea except Mooren's ulcer which is ulcerative keratitis.

---

**QUESTION 3**

A pregnant woman comes with weakness and elevated liver function tests. She is in her 35th week of pregnancy. The prothrombin time is normal. The smear of blood shows fragmented red cells. Platelet count is low.

What is the treatment?

- A. Delivery of the baby
- B. Fresh frozen plasma
- C. Prednisone

Correct Answer: A

Deliver the baby with the HELLP syndrome. HELLP syndrome stands for hemolysis, elevated liver function tests, and low platelets. This disorder is idiopathic and can be distinguished from DIC by the normal coagulation studies, such as the prothrombin time and aPTT.

---

**QUESTION 4**

19-year-old sexually active lady came for her annual check-up, she is otherwise healthy using no contraceptive, her pap smear and all investigations are normal.

What will you suggest regarding her next check-up?

- A. After 6 months
- B. After 5 years
- C. After 1 year
- D. After 3 years

Correct Answer: D

Usually pap smear should be started after 21 years of age regardless of sexual activeness but if a female is sexually active and u suspect any abnormality then u can do pap smear before 21 years of age as in this case. Pap Smear = Negative, then should be followed for screening after 3 years. Pap smear + HPV Testing = Negative, then should be followed after 5 years

---

**QUESTION 5**

A male presented with headache, tinnitus and nausea thinking that he has a brain tumor. He had just secured a job in a prestigious company and he thinks that he might not meet its standards. CNS exam, CT, all within normal.

What is the diagnosis?

- A. Conversion reaction
- B. Generalized Anxiety disorder
- C. Panic attack
- D. Hypochondriasis

Correct Answer: D

Diagnostic criteria for hypochondriasis include the following: The patient has a preoccupying fear of having a serious disease. The preoccupation persists despite appropriate medical evaluation and reassurance. The belief is not of delusional intensity (as in delusional disorder, somatic type) and is not Restricted to a concern about appearance (as in persons with BDD). The preoccupation causes clinically significant distress or impairment.